


AD-A184 259 PRODUCTION AND SELECTION OF MONOCLONAL ANTIBODIES FOR 1/1
DETECTION OF SMALL (U) CALIFORNIA UNIV OAKLAND NAVAL
BIOSCIENCES LAB A E KARU 1987 UC-NBL-964
UNCLASSIFIED N00014-81-C-0570 F/G 6/5 NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

SECURITY

AD-A184 259

DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS NONE	
2a. SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE N/A		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NBL No. 964		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6a. NAME OF PERFORMING ORGANIZATION University of California	6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) Code 1141 800 North Quincy Ave Arlington, VA 22217-5000	
6c. ADDRESS (City, State, and ZIP Code) Naval Biosciences Laboratory Naval Supply Center Oakland, California 94625	8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-81-C-0570
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Avenue Arlington, Va 22217-5000	10. SOURCE OF FUNDING NUMBERS		
	PROGRAM ELEMENT NO 61153N	PROJECT NO RR041-05	TASK NO RR041-05-03 WORK UNIT ACCESSION NO NR204-123
11. TITLE (Include Security Classification) (U) PRODUCTION AND SELECTION OF MONOCLONAL ANTIBODIES FOR DETECTION OF SMALL MOLECULAR WEIGHT COMPOUNDS			
12. PERSONAL AUTHOR(S) Karu, Alexander E.			
13a. TYPE OF REPORT Summary Report	13b. TIME COVERED FROM 861001 TO 870930	14. DATE OF REPORT (Year, Month, Day) 1987	15. PAGE COUNT 1
16. SUPPLEMENTARY NOTATION 101 Annual Meeting Association of Official Analytical Chemists, S.F., CA Sept 1987.			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
06	03		
		Keywords: Monoclonal antibodies, Anti-hapten	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Production of useful anti-hapten monoclonal antibodies required integrating carefully chosen immunization, selection, screening and cell management techniques.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Head, Biological Sciences Div, ONR		22b. TELEPHONE (Include Area Code) (202) 698-4986	
		22c. OFFICE SYMBOL ONR Code 1141	

DTIC
SELECTED
SEP 09 1987
D

20/22

964



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Invited talk, Symposium on Biotechnology, 101st Annual Meeting, Association of Official Analytical Chemists, San Francisco CA, September 15, 1987

ALEXANDER E. KARU (Hybridoma Center, College of Natural Resources, Univ. of Calif., Berkeley, CA 94720) : Production and Selection of Monoclonal Antibodies for Detection of Small Molecular Weight Compounds.

Production of useful anti-hapten monoclonal antibodies requires integrating carefully chosen immunization, selection, screening, and cell management techniques. This talk will survey principles and methods that many laboratories use but few systematically evaluate. Design and hapten density of the conjugates used for immunizing and screening are critical for successful hybridoma production. Immunization protocols should be set up to avoid causing suppression of hapten-specific responses. Unrelated mouse strains should be immunized because various inbred lines express different immunoglobulin variable-region genes, and other immunoregulatory genes affect response to hapten. Adjuvants can be chosen with respect to their behavior as immunostimulators, solvents, and depots for hapten-carrier complexes. Immune response modifiers may be used to stimulate helper T-cell responses or inhibit suppressor T-cell responses to the hapten, or to promote antibody production in B cells. Use of conditioned media instead of "feeder" cells simplifies hybridoma culture and allows growth at low cell densities. This maximizes detectability and minimizes loss of good clones. The best hybridomas will be very rare, and may differ greatly in affinity and specificity for the immunizing hapten and its analogs. Automated sampling systems and computer-assisted screening are essential for rapid identification of the most useful cell lines. Supported by Office of Naval Research Contract N00014-81-C-0570 and NSF Grant DMB-86-12088.

END

10-87

DTIC